arXiv:1304.4406v1 [physics.soc-ph] 16 Apr 2013

Cities as nuclei of sustainability?

Diego Rybski[a,*], Till Sterzel[a], Dominik E. Reusser[a], Christina Fichtner[a],
Jürgen P. Kropp[a,b]

[a] *Potsdam Institute for Climate Impact Research, 14412 Potsdam, Germany, EU*
[b] *Institute of Earth and Environmental Science, University of Potsdam, 14476 Potsdam, Germany, EU*

Abstract

Analyzing CO_2 emission inventories of 256 cities from 33 countries we find power-law correlations between the emissions and city size, measured in population. The results suggest that in developing countries more CO_2 per capita is emitted in large cities, i.e. they tend to exhibit super-linear correlations and doubling the population of any city implies up to 110% increase of emissions. For developed countries the results suggest the opposite, i.e. linear or sub-linear correlations, implying better efficiency of large cities, doubling the population of any city implies only 80% increase of emissions. The transition occurs at approx. $10,000$\$ GDP/cap. We derive how the total emissions of an entire country relate with the power-law correlations and find that the size of the most populated city is dominating in the case of linear and super-linear correlations. The size of the largest city has no influence in the case of sub-linear correlations. We conclude that from the climate change mitigation point of view, urbanization is desirable in developed countries and should be accompanied by efficiency increasing mechanisms in developing countries. More data acquisition is needed to support our empirical findings.

Keywords: CO_2 emissions, Cities, Urban Area, Scaling, Development, Transition

---

Journal:

---

[*] ca-dr@rybski.de

1. Introduction

Cities are the place where most people live (UNPD, 2007) and the largest part of current emissions is associated to urban areas (Hoornweg et al., 2011). However, a generalization of the dirty and pollutant metropolis may be wrong considering the finding that various positive features, such as innovation or GDP, are more effective in large (we always refer to population when we speak about the size of a city) cities compared to smaller ones (Bettencourt and West, 2010). Accordingly, it is of interest how city size and CO_2 emissions are related, i.e. to which extent a similar synergetic effect can be found. The outcome could shed light on the question about sustainability of urban life (Bettencourt and West, 2010).

While Dodman (2009) argues that per capita emissions from cities are lower than the corresponding country value, Hoornweg et al. (2011) find that emissions on different scales vary considerably for the same resident depending on whether production- or consumption-based figures are considered. Brown et al. (2009) conclude that metropolitan areas can have large impact towards a climate friendly environment.

Usually, proportionality between city size and city emissions is assumed, i.e. the more inhabitants, the more CO_2 is emitted. However, this assumption has neither been verified nor disproved empirically, which is mostly due to the lack of city scale emission values. In case the linearity does hold, it could be used to estimate emissions of cities for which no data is available. Currently, linearity is assumed for downscaling emissions from country to high resolution raster scale, e.g. (Rayner et al., 2010; Oda and Maksyutov, 2011). In case the correlations between population and emissions are non-linear, the question emerges which form they follow and which factors determine their shape. Particularly interesting is the influence of the countries development. Since the standard procedure of emission reporting to UNFCCC is well established (IPCC, 2006), the analogous decomposition into emissions from cities could help to assess their sustainability, i.e. a *reporting scheme for cities* is needed.

We investigate CO_2 emissions versus city size of 256 large cities situated in 33 different countries. We relate the two quantities with a power-law, using the exponent γ to characterize whether the emissions depend linearly, super- or sub-linearly on the city size, i.e. whether large cities emit more CO_2 emissions per capita compared to small cities, or vice versa. In order to examine the influence of economic development, we check how the exponent of the power-law relates to the GDP per capita of the countries. We find a power-law relation between city size and emissions, whereas the exponent depends on the development of the country. Finally, we derive how the total emissions of a country depend on the size of the largest city. We conclude that a transition occurs at $\gamma = 1$ from super- to sub-linear behavior, where the domination of the largest city is overcome.

2. Data

CO_2 emission figures (we have used CO_2 *equivalent* figures and use the notation CO_2 for simplicity) were retrieved from collections of urban GHG emission inventories published in peer reviewed journals or reports from research institutes and non-governmental organizations. From the plethora of figures available we compiled a set of 256 per capita urban GHG emission figures from a variety of different GHGs methodologies, sources, and sectors. The emissions accounted for are converted to CO_2 if necessary. 231 inventories (90%) explicitly include CO_2 emissions albeit from varying sectors and sources. 25 (10%) inventories were unspecific about which emissions were accounted for. 136 (53%) inventories account for CO_2 emissions, and a further 10 (4%) are ambiguous as to whether they account for additional CH_4 and N_2O emissions 82 (32%) inventories account for all emissions covered by the Kyoto Protocol (CO_2, CH_4, N_2O, SF_6, HFCs and PFCs), and 3 (1%) inventories account for CO_2, CH_4, N_2O only.

Published figures were largely drawn from primary sources such as city inventories covering emissions in their territories. If necessary, figures were traced back to their original publication source for verification. The per capita emissions considered range from 0.1 t/cap for Rajshahi, Bangladesh ICLEI (2009) and 43.7 t/cap for Aberystwyth, Wales (derived from Dore et al. (2006) and Carbon Trust (2007/08)). For 23 cities both city and metropolitan region figures available were considered as they can diverge substantially (Hoornweg et al., 2011).

Publications by Brown et al. (2008), Brown et al. (2009), Hoornweg et al. (2011), Kennedy et al. (2009), and Sovacool and Brown (2010) Carbon Trust (2007/08), Dore et al. (2006), and ICLEI (2009) account for 248 of the inventory figures (93%).

A proposal for a principal standard for urban emission inventories has recently been suggested by C40 Cities Climate Leadership Group and ICLEI Local Governments for Sustainability in collaboration with World Resources Institute, World Bank, UNEP, and UN-HABITAT (C40 and ICLEI, 2012).

However, for currently existing inventories a multitude of methodologies have been and are still in use, ranging widely in terms of the area, sector or emission(s) of interest, the precision of the methodologies applied, and in how far the inventories abided by the methodology and traceably documented the input, methods and findings in the published results. The literature and its urban emissions figures considered for this study reflect this heterogeneity which could be a source of dispersion.

3. City emissions

We group our database of 256 city emission inventories according to the country's GDP/cap, striving for a balanced number of cases in each of the five groups. Based on these groups, we begin our analysis by plotting the annual amount of CO_2 emission (We always refer to *annual* figures and omit "annual" for simplicity) e against the corresponding city population s in Fig. 1. In order to make the city emissions of different countries comparable, we divide by the emissions per capita of the corresponding countries. We find correlations which approximately follow power-laws according to

$$e = a\, s^{\gamma}\,, \tag{1}$$

where a is a proportionality constant – for the linear case ($\gamma = 1$) it is identical to the emissions per capita. The exponent γ determines how strongly the emissions increase with city size: $\gamma = 1$ (linear) the larger the population, the more emissions; $\gamma < 1$ (sub-linear) large cities emit less CO_2 than expected from emissions per capita of small cities; and $\gamma > 1$ (super-linear) large cities emit more CO_2 compared to small ones. As mentioned above, similar power-laws have been found for other quantities, i.e. crime, GDP, income, and patents (Bettencourt and West, 2010). In addition to the evaluation by the five country groups, we also estimated γ for the three countries with more than 10 city inventories in our database: India (part of Group 2), UK (part of Group 4) and US (coincident with Group 5).

The obtained exponents γ of each group and country are plotted against the corresponding GDP per capita in Fig. 2. In the case of groups with more than one country, a weighted country GDP was calculated based on the number of occurrences of cities from each country in the group. The Figure suggests that countries with low GDP/cap values tend to exhibit $\gamma > 1$ and countries with larger GDP/cap values tend to exhibit smaller γ, around or



Figure 1: City efficiency in terms of CO_2 emissions. Per capita (annual) CO_2 emissions versus population and regressions according to Eq. (1). The emission values have been normalized by dividing through the country emissions per capita. City emission figures have been sorted and separated into groups according to the GDP/cap of the countries. (a) Group 1 ($R^2 = 0.65$): Philippines (1), Bangladesh (4), Nepal (3). (b) Group 2 ($R^2 = 0.78$): China (5), South Africa (1), Thailand (2), Greece (1), Sri Lanka (4), Slowenia (1), Czech Republic (1), Portugal (1), Spain (2), Mexico (1), South Korea (2), Indonesia (1), Brazil (3), India (42), Bhutan (2) and India (42, brown open triangles). (c) Group 3 ($R^2 = 0.79$): Germany (4), Italy (4), Singapore (1), Belgium (1), Finland (1), Japan (1), France (1), Sweden (1). (d) Group 4 ($R^2 = 0.75$): UK (35), Netherlands (1), Canada (4), Australia (1), Switzerland (1), Norway (1) and UK (35, grey open squares). (e) Group 5 ($R^2 = 0.92$): USA (122).



Figure 2: Economic development and city efficiency in terms of CO_2 emissions. Determined exponents γ from Eq. (1) versus the corresponding GDP per capita. For the same groups and countries as in Fig. 1 the obtained power-law exponents γ are plotted against the weighted average of the country's GDP/cap.

below 1. Notwithstanding the weak evidence, the question emerges if there is a difference in the scaling of developing and developed countries, i.e. if cities of (economically) developing countries tend to a super-linear relation between population and CO_2 emissions ($\gamma > 1$), while cities of (economically) developed countries tend to a sub-linear relation ($\gamma < 1$). This would imply a transition at $\gamma \simeq 1$. In other words, in economically stronger countries, large cities tend to be more *efficient* and in economically weaker countries large cities tend to be more *inefficient* in terms of CO_2 emissions per capita.

4. Country emissions

Relating emissions from cities to the emissions from an entire country could help to understand the drivers of emissions. In order to estimate the emissions of an entire country from the above relation between city size and emissions, it is necessary to take into account how many cities exist of each size including towns and villages. Such a relation characterizing the size distribution of cities is Zipf's law (Zipf, 2012; Auerbach, 1913) which is based on empirical data and expresses that city sizes follow broad distributions, i.e. the probability density $p(s) \sim s^{-\zeta}$. For cities it has been found that the Zipf exponent is $\zeta \simeq 2$, see e.g. (Rozenfeld et al., 2011) and references therein. In addition to the assumptions over the size distribution of cities, we can assume that emissions from non-urban regions are negligible.

Under this assumption, the total emissions of a country are given by the sum over the number of cities of certain size times the typical emissions of the corresponding city size. Assuming $\zeta \simeq 2$, we obtain E, the expression for the emissions from the considered country,

$$E \simeq \begin{cases} \frac{a}{\gamma-1} s_{\max}^{\gamma-1} & \text{if } \gamma > 1 \\ a \ln(s_{\max}) & \text{if } \gamma \simeq 1 \\ \frac{a}{1-\gamma} & \text{if } \gamma < 1 \end{cases}, \tag{2}$$

see Appendix A.

The three distinct cases are illustrated in Fig. 3 (in semi-logarithmic representation):

$\gamma > 1$: The country emissions E as a function of the size of the largest city $\ln(s_{\max})$ have a concave shape.



Figure 3: Illustration of country emissions as a function of the population of the largest city, $s_{\max}$. The curves are given by Eq. (A.4) and (A.5) with $\zeta = 2$ and constant a.

$\gamma \simeq 1$: E as a function of $\ln(s_{\max})$ is linear. E increases logarithmically with $s_{\max}$, independent of γ.

$\gamma < 1$: E as a function of $\ln(s_{\max})$ has a convex shape. The emissions of a country are independent of the population of the largest city for sufficiently large $s_{\max}$ and approach an asymptotic constant value.

In any case, the proportionality constant a remains. The exponent γ reflects the development of the country and the special case $\gamma = 1$ marks a transition performed by the countries while economically growing.

5. Discussion

The large number of people living and working in cities implies frequent human interaction, which in turn can have positive or negative facets. In the case of CO_2 emissions a differentiated view is necessary. Depending on the development of the considered countries different processes seem to be responsible for more or less emissions.

On a country scale, it was found that per capita CO_2 emissions correlate exponentially with the development as measured by the human development index (HDI) (Costa et al., 2011). Per capita CO_2 emissions are determined by various factors. In addition to the type of energy sources and efficiency of devices, livelihood conditions and lifestyles play an important role. Out of 16 elements used to describe livelihood conditions, the six elements water availability, nutrition, energy supply, housing/shelter, sanitation and social cohesion where found to be related to significant CO_2 emissions, while the other appear only loosely coupled to CO_2 emissions (Reusser et al., 2013). Emissions are determined by a combination of advancement of technology and level of consumption. Thus, people in developed countries cause more emissions – despite advanced technology – simply due to their increased consumption of all kinds of resources and services. Per capita emissions are much higher in developed countries and adjusting for emissions embodied in international trade further increases this discrepancy (Caldeira and Davis, 2011; Peters and Hertwich, 2008).

We do not know the reasons for the suggested γ-transition but the processes seem to depend on the development of the considered country. A possible explanation could be the allocation of facilties (Gastner and Newman, 2006). Um et al. (2009) find that commercial facilities exhibit linear and public ones sub-linear scaling with population. If one assumes that the facility types differ by their emissions and that the development of the public sector is more or less pronounced, then different emissions scaling can be expected depending on the development of the countries.

Another explanation could be a different structure of the city's economy shifting from industry to more services, which has been discussed as an important mechanism of carbon reduction due to modernization (Lankao et al., 2008). Such differences in the structure and change of the economies dependent on the degree of development and are evident at least on the scale of countries (Lutz et al., 2013). The to-date data availability is insufficient to test if the structure of the city's economy is changing differently depending on the level of development.

In addition to the above mentioned lifestyles and economic composition, the influence of city size can have positive or negative effects in terms of CO_2 emissions. On the one hand, the compactness of cities implies shorter distances and accordingly less emissions from transport. While generally the transport system is better established in developed countries, the distribution of developed and developing countries is inhomogeneous regarding the climate. On the other hand, the urban heat island effect is more pronounced in large cities but might depend on the climate where they are located (Zhou et al., 2013), which in turn can affect the heating and cooling behavior of people, see e.g. (Olonscheck et al., 2011). These are only two aspect which could lead to super- or sub-linear scaling depending on the development of the considered countries.

Unfortunately, we can only provide poor evidence of a development dependent γ. The analyzed figures have been collected by various authors and we cannot exclude differences in the technical details of the inventories. This shows that the available data on city emissions is still insufficient and we urge a systematic reporting of CO_2 emissions from cities, for example as recently proposed (C40 and ICLEI, 2012). We plan to verify our results in future research based on updated city inventories as they become available. The same holds for alternative data sources, such as the high resolution fossil fuel combustion CO_2 emissions fluxes for the US (Gurney et al., 2009) or data from the Greenhouse gases Observing SATellite (GOSAT) (Hammerling et al., 2012) and local detailed GHG monitoring programmes (e.g. Tollefson, 2012).

Acknowledgments

We thank Anika Nockert and Stephan Huber for useful discussions. We appreciate financial support by the Potsdam Research Cluster for Georisk Analysis, Environmental Change and Sustainability (PROGRESS) and the RAMSES project (FP7).

Appendix A. Derivation of Eq. (2)

According to Zipf's law the city sizes follow the probability density

$$p(s) = c\, s^{-\zeta} \qquad (1 \leq s \leq s_{\max})\,, \tag{A.1}$$

where ζ is the Zipf exponent and c is a constant which is determined from normalization,

$$c = \frac{\zeta - 1}{1 - s_{\max}^{1-\zeta}}\,, \tag{A.2}$$

and $s_{\max}$ is the population of the largest city.

The total emissions of a country are given by the integral over the product of probability density of city sizes, Eq. (A.1), and the emissions of the considered city sizes, Eq. (1),

$$E \;=\; \int p(s)e(s)\mathrm{d}s \;=\; \int c\, s^{-\zeta}\, a\, s^{\gamma}\mathrm{d}s\,, \tag{A.3}$$

where E are the total emissions of a country.

Using Eq. (A.2) and putting factors outside the integral gives:

$$E = \frac{a(\zeta-1)}{1-s_{\max}^{1-\zeta}} \int_1^{s_{\max}} s^{\gamma-\zeta} \mathrm{d}s \,.$$

In order to integrate, one needs to distinguish two cases.

First, $\gamma = \zeta - 1$ leads to

$$E = \frac{a(\zeta-1)\ln(s_{\max})}{1-s_{\max}^{1-\zeta}} \,. \tag{A.4}$$

This means, the country emissions are determined by the constants a and ζ as well as depend on the size of the largest city $s_{\max}$. Second, $\gamma \neq \zeta - 1$ leads to

$$E = \frac{a(\zeta-1)}{\gamma-\zeta+1} \frac{s_{\max}^{\gamma-\zeta+1}-1}{1-s_{\max}^{1-\zeta}} \,. \tag{A.5}$$

Again, E is determined by the constants a and ζ, but in this case there is a power-law term of $s_{\max}$ instead of the logarithmic one.

Equations (A.4) and (A.5) can be considered in the limit $s_{\max} \gg 1$ leading to three different cases

$$E = \begin{cases} \frac{a(\zeta-1)}{\gamma-\zeta+1} s_{\max}^{\gamma-\zeta+1} & \text{if } \gamma > \zeta - 1 \\ a(\zeta-1)\ln(s_{\max}) & \text{if } \gamma = \zeta - 1 \\ \frac{a(\zeta-1)}{\zeta-\gamma-1} & \text{if } \gamma < \zeta - 1 \end{cases} \,. \tag{A.6}$$

For $\zeta > 1$, and if we additionally assume $\zeta \simeq 2$, we obtain

$$E \simeq \begin{cases} \frac{a}{\gamma-1} s_{\max}^{\gamma-1} & \text{if } \gamma > 1 \\ a\ln(s_{\max}) & \text{if } \gamma \simeq 1 \\ \frac{a}{1-\gamma} & \text{if } \gamma < 1 \end{cases} \,. \tag{A.7}$$

References

Auerbach, F., 1913. Das Gesetz der Bevölkerungskonzentration. Petermanns Geogr. Mitteilungen 59 (74), 73–76.

Bettencourt, L., West, G., 2010. A unified theory of urban living. Nature 467 (7318), 912–913.

Brown, M. A., Southworth, F., Sarzynski, A., 2008. Shrinking the carbon footprint of metropolitan america. Tech. rep.

Brown, M. A., Southworth, F., Sarzynski, A., 2009. The geography of metropolitan carbon footprints. Policy and Society 27 (4), 285–304.

C40, ICLEI, 2012. Global Protocol for Community-Scale Greenhouse Gas Emissions. Tech. Rep. May, C40 Cities Climate Leadership Group and ICLEI Local Governments for Sustainability in collaboration with: World Resources Institute, World Bank, UNEP, and UN-HABITAT, São Paulo.

Caldeira, K., Davis, S. J., 2011. Accounting for carbon dioxide emissions: A matter of time. Proc. Nat. Acad. Sci. U.S.A. 108 (21), 8533–8534.

Carbon Trust, 2007/08. Annual report. Tech. rep., http://reports.investis.com/reports/cbnt_ar_2008_en/report.php.

Costa, L., Rybski, D., Kropp, J. P., 2011. A human development framework for CO_2 reductions. PLoS ONE 6 (12), e29262.

Dodman, D., 2009. Blaming cities for climate change? An analysis of urban greenhouse gas emissions inventories. Environ. Urban. 12 (1), 185–201.

Dore, C. J., Watterson, J. D., Murrells, T. P., Passant, N. R., Hobson, M. M., Baggott, S. L., Thistlethwaite, G., Goodwin, J. W. L., King, K. R., Adams, M., Walker, C., Downes, M. K., Coleman, P. J., Stewart, R. A., Wagner, A., Sturman, J., Conolly, C., Lawrence, H., Li, Y., Jackson, J., Bush, T., Grice, S., Brophy, N., 2006. Uk emissions of air pollutants 1970 to 2004. Tech. Rep. December, National Environmental Technology Centre.

Gastner, M. T., Newman, M. E. J., 2006. Optimal design of spatial distribution networks. Phys. Rev. E 74 (1), 016117.

Gurney, K. R., Mendoza, D. L., Zhou, Y. Y., Fischer, M. L., Miller, C. C., Geethakumar, S., Du Can, S. D., 2009. High resolution fossil fuel combustion CO_2 emission fluxes for the united states. Environ. Sci. Technol. 43 (14), 5535–5541.

Hammerling, D. M., Michalak, A. M., O'Dell, C., Kawa, S. R., Apr. 2012. Global CO 2 distributions over land from the Greenhouse Gases Observing Satellite (GOSAT). Geophys. Res. Lett. 39 (8).

Hoornweg, D., Sugar, L., Gomez, C. L. T., 2011. Cities and greenhouse gas emissions: moving forward. Environ. Urban. 23 (1), 207–227.

ICLEI, 2009. Energy and carbon emissions profiles of 54 south asian cities. Tech. rep., ICLEI - Local Governments for Sustainability, New Delhi.

IPCC, 2006. 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Prepared by the National Greenhouse Gas Inventories Programme. IGES, Japan.

Kennedy, C. A., Ramaswami, A., Carney, S., Dhakal, S., 2009. Greenhouse gas emission baselines for global cities and metropolitan regions. Tech. rep., http://siteresources.worldbank.org/INTURBANDEVELOPMENT/Resources/336387-12565668 1268260567624/KennedyComm.pdf.

Lankao, P. R., Nychka, D., Tribbia, J. L., 2008. Development and greenhouse gas emissions deviate from the ’modernization’ theory and ’convergence’ hypothesis. Clim. Res. 38 (1), 17–29.

Lutz, R., Spies, M., Reusser, D. E., Kropp, J. P., Rybski, D., 2013. Characterising the development of sectoral gross domestic product composition. submitted.

Oda, T., Maksyutov, S., 2011. A very high-resolution (1 km x 1 km) global fossil fuel CO_2 emission inventory derived using a point source database and satellite observations of nighttime lights. Atmos. Chem. Phys. 11 (2), 543–556.

Olonscheck, M., Holsten, A., Kropp, J. P., 2011. Heating and cooling energy demand and related emissions of the German residential building stock under climate change. Energy Policy 39 (9), 4795–4806.

Peters, G. P., Hertwich, E. G., 2008. CO2 embodied in international trade with implications for global climate policy. Environ. Sci. Technol. 42 (5), 1401–1407.

Rayner, P. J., Raupach, M. R., Paget, M., Peylin, P., Koffi, E., 2010. A new global gridded data set of CO_2 emissions from fossil fuel combustion: Methodology and evaluation. J. Geophys. Res.-Atmos. 115, D19306.

Reusser, D. E., Costa, L., Lissner, T., Pradhan, P., Holsten, A., Rybski, D., Reusswig, F., Kropp, J. P., 2013. Relating climate compatible development and human livelihood. in preparation.

Rozenfeld, H. D., Rybski, D., Gabaix, X., Makse, H. A., 2011. The area and population of cities: New insights from a different perspective on cities. Am. Econ. Rev. 101 (5), 2205–2225.

Sovacool, B. K., Brown, M. A., 2010. Twelve metropolitan carbon footprints: A preliminary comparative global assessment. Energy Policy 38 (9), 4856–4869.

Tollefson, J., 2012. Megacities move to track emissions. Nature 492 (7427), 20–21.

Um, J., Son, S. W., Lee, S. I., Jeong, H., Kim, B. J., 2009. Scaling laws between population and facility densities. Proc. Nat. Acad. Sci. U.S.A. 106 (34), 14236–14240.

UNPD, 2007. World urbanization prospects: The 2007 revision. Tech. rep., united Nations Population Division, New York.

Zhou, B., Rybski, D., Kropp, J. P., 2013. Urban heat island - A study based on a vast number of urban agglomerations. in preparation.

Zipf, G. K., 2012. Human Behavior and the Principle of Least Effort: An Introduction to Human Ecology (Reprint of 1949 Edition). Martino Publishing, Manfield Centre, CT.